

**TABLE 1-5. United States Rates of Low Birthweight, and Measures of Prenatal Care, 1980-2000**

Year	Low Birthweight	First Trimester Care	No Care	Inadequate Care	Third Trimester Care	Less Than Five Visits
1980	68.4	763.6	13.5	87.2	38.1	69.4
1981	68.1	763.5	14.1	87.1	38.4	68.6
1982	67.5	759.3	15.9	90.8	39.9	71.9
1983	68.2	760.6	17.0	88.7	39.7	69.9
1984	67.2	764.5	17.1	87.8	39.4	68.7
1985	67.5	763.1	17.0	88.0	40.6	67.6
1986	68.1	760.4	19.3	89.6	41.1	68.4
1987	69.0	760.0	20.1	90.5	41.8	68.8
1988	69.3	760.5	18.8	90.4	42.1	68.4
1989	70.5	754.5	21.8	96.3	42.7	74.6
1990	69.7	758.3	19.8	91.3	41.1	70.4
1991	71.2	762.5	19.1	86.7	38.6	66.6
1992	70.8	777.5	17.3	78.6	34.5	60.6
1993	72.2	789.0	16.0	72.7	32.4	55.2
1994	72.8	802.2	13.6	66.9	30.4	50.4
1995	73.2	812.7	12.3	63.0	30.2	46.7
1996	73.9	818.6	11.8	60.5	28.2	44.7
1997	75.1	825.3	12.2	58.1	27.0	44.5
1998	76.0	828.3	11.9	57.9	27.0	44.1
1999	76.0	832.0	11.2	56.3	25.8	43.4
2000*	76.0	832.0	not available	not available	not available	not available

Inadequate prenatal care is defined as care that began in the third trimester or consisted of less than five visits.

All rates per 1,000 live births. Rates and percentages are calculated excluding missing and unknown values.

\* Martin JA, Hamilton BE, Ventura SJ. Births: Preliminary data for 2000. National vital statistics reports; vol 49 no 5. Hyattsville, Maryland: National Center for Health Statistics. 2001.

**TABLE 1-6. Oregon Rates of Low Birthweight, and Measures of Prenatal Care, 1980-2000**

Year	Low Birthweight	First Trimester	No Care	Inadequate Care	Third Trimester Care	Less than Five Visits
1980	50.4	780.8	5.5	58.0	35.2	41.4
1981	48.5	775.6	8.9	63.1	38.6	43.0
1982	49.2	769.3	11.2	70.3	41.0	48.0
1983	50.0	775.3	11.3	66.5	38.5	44.9
1984	51.5	771.5	11.0	68.2	41.1	46.2
1985	51.3	752.0	12.1	72.9	43.7	47.5
1986	51.3	738.7	11.7	83.3	52.1	54.6
1987	54.0	736.8	16.5	86.2	50.3	58.5
1988	52.6	738.8	13.8	83.6	49.9	54.7
1989	52.2	750.7	12.0	73.2	42.9	48.7
1990	50.1	757.1	10.7	70.0	43.4	45.1
1991	49.2	768.2	8.7	61.0	37.4	38.6
1992	51.8	787.0	8.2	52.6	31.4	34.0
1993	52.5	794.6	7.6	51.7	30.4	33.8
1994	53.0	790.9	8.5	57.8	34.3	36.4
1995	54.9	787.7	8.6	58.4	34.7	38.2
1996	53.5	799.3	7.1	53.7	31.7	34.8
1997	55.0	811.2	6.7	50.0	29.6	32.3
1998	53.7	807.2	7.2	53.5	30.7	35.3
1999	53.9	809.9	7.3	53.7	29.6	35.7
2000	56.6	812.8	8.5	55.9	29.8	36.6

Inadequate prenatal care is defined as care that began in the third trimester or consisted of less than five visits.

All rates are per 1,000 live births.

Rates and percentages are calculated excluding missing and unknown values.