

**TABLE 6-20. Tobacco-linked Deaths by Cause of Death, Oregon Residents, 2007**

Selected Causes of Death (and their ICD-10 codes)	Total	Linked <sup>1</sup>		Not Linked		Unknown	
		Number	Percent	Number	Percent	Number	Percent
Total <sup>2</sup> .....	31,433	6,965	22.2	16,737	53.2	7,731	24.6
Malignant Neoplasms .....	3,410	1,922	56.4	957	28.1	531	15.6
Oral cavity, lip, pharynx (C00.0-C14.8) .....	94	51	54.3	30	31.9	13	13.8
Esophagus (C15) .....	205	83	40.5	69	33.7	53	25.9
Stomach (C16) .....	117	19	16.2	67	57.3	31	26.5
Pancreas (C25) .....	473	43	9.1	332	70.2	98	20.7
Larynx (C32) .....	29	15	51.7	8	27.6	6	20.7
Lung, bronchi, and trachea (C33-C34) .....	2,032	1,633	80.4	180	8.9	219	10.8
Cervix uteri (C53) .....	31	2	6.5	22	71.0	7	22.6
Kidney, other urinary tract (C64-C65) .....	136	14	10.3	91	66.9	31	22.8
Urinary bladder (C67) .....	191	60	31.4	76	39.8	55	28.8
Acute Myeloid Leukemia (C92.0) .....	102	2	2.0	82	80.4	18	17.6
Cardiovascular Disease .....	8,588	1,850	21.5	4,180	48.7	2,558	29.8
Ischemic heart disease (I20-I25) .....	3,935	1,119	28.4	1,593	40.5	1,223	31.1
Other heart disease (I00-I09, I26-I51) .....	2,426	336	13.8	1,401	57.7	689	28.4
Cerebrovascular disease (I60-I69) .....	1,833	258	14.1	1,023	55.8	552	30.1
Atherosclerosis (I70) .....	124	30	24.2	65	52.4	29	23.4
Aortic aneurysm (I71) .....	168	70	41.7	58	34.5	40	23.8
Other arterial disease (I72-I78) .....	102	37	36.3	40	39.2	25	24.5
Respiratory Diseases .....	2,290	1,553	67.8	419	18.3	318	13.9
Pneumonia and influenza (J10-J18) .....	481	57	11.9	299	62.2	125	26.0
Bronchitis and emphysema (J40-J43) .....	253	218	86.2	15	5.9	20	7.9
Other chronic airways obstruction (J44) .....	1,556	1,278	82.1	105	6.7	173	11.1
Perinatal Conditions <sup>3</sup> .....	93	1	1.1	84	90.3	8	8.6
Selected Perinatal Conditions <sup>4</sup> .....	55	—	—	52	94.5	3	5.5
Sudden Infant Death Syndrome (R95) .....	38	1	2.6	32	84.2	5	13.2
Other causes .....	17,052	1,639	9.6	11,097	65.1	4,316	25.3

<sup>1</sup> The Oregon death certificate asks 'Did tobacco use contribute to death?' followed by four checkboxes: 'Yes,' 'No,' 'Probably,' and 'Unknown.' The linked category includes deaths listed as 'Yes' or 'Probably.'

<sup>2</sup> The causes of death shown in this table are those linked to tobacco use by the federal Centers for Disease Control and Prevention (CDC). Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses -- United States, 1997-2001. MMWR 2005; 54:625-628.).

<sup>3</sup> The number of infant deaths resulting from exposure to tobacco combustion products is underreported.

<sup>4</sup> The category includes the following conditions: other disorders related to short gestation and low birthweight (P07), respiratory distress of newborn (P22), congenital pneumonia (P23), neonatal aspiration syndromes (P24), and other respiratory conditions originating in the perinatal period (P25-P28).

— Quantity is zero.