

TABLE 6-53. Crude Death Rates for Selected Leading Causes of Mortality, United States, 1994-2008

Year	Total	Heart Disease	Cancer	CLRD ¹	Cerebrovascular Disease	Unintended Injury	Alzheimer's Disease	Diabetes	Pneumonia and Influenza
1994	875.4	277.1	207.1	40.6	61.9	36.0	12.3	22.2	21.8
1995	880.0	276.5	206.8	40.8	63.1	36.4	13.3	23.0	22.3
1996	872.5	272.4	205.3	41.6	63.3	36.7	13.4	23.7	22.3
1997	864.7	267.6	203.5	42.4	62.7	36.6	13.8	23.9	22.5
1998	864.2	263.7	202.1	43.1	58.9	37.1	13.8	24.4	23.7
1999	877.0	265.9	201.6	45.5	61.4	35.9	16.3	25.1	23.4
2000	873.6	257.9	200.5	44.9	60.3	34.0	17.8	24.9	24.3
2001	848.5	245.8	196.0	43.7	57.9	35.7	18.9	25.1	22.0
2002	847.3	241.7	193.2	43.3	56.4	37.0	20.4	25.4	22.8
2003	841.9	235.6	191.5	43.5	54.2	37.6	21.8	25.5	22.4
2004	816.5	222.2	188.6	41.5	51.1	38.1	22.5	24.9	20.3
2005	825.9	220.0	188.7	44.2	48.4	39.7	24.2	25.3	21.3
2006	810.4	211.0	187.0	41.6	45.8	40.6	24.2	24.2	18.8
2007	803.6	204.3	186.6	42.4	45.1	41.0	24.7	23.7	17.5
2008	813.0	202.9	186.0	46.4	44.1	40.1	27.1	23.2	18.5

Year	Suicide	Hyper-tension	Alcohol-induced ³	Parkin-son's Disease	Homicide (excluding legal intervention)	Acquired Immune Deficiency Syndrome	Congenital Anomalies	Arterio-sclerosis ²	Amyo-trophic Lateral Sclerosis
1994	12.0	5.0	7.6	3.8	9.4	17.5	4.1	6.4	1.5
1995	11.9	5.2	7.6	4.1	8.6	17.7	4.1	6.2	1.5
1996	11.6	5.5	7.3	4.5	7.8	12.7	4.0	6.1	1.6
1997	11.4	5.7	7.2	4.6	7.3	6.7	3.9	5.8	1.6
1998	11.3	5.9	7.1	4.9	6.6	5.4	3.9	5.5	1.6
1999	10.7	6.2	7.1	5.4	6.2	5.4	3.8	5.5	1.9
2000	10.3	6.5	7.0	5.7	5.9	5.2	3.8	5.2	1.9
2001	10.8	6.8	7.0	5.8	7.1	5.0	3.7	4.9	1.9
2002	11.0	7.0	6.9	5.9	6.1	4.9	3.7	4.8	2.0
2003	10.8	7.5	7.0	6.2	6.1	4.7	3.6	4.5	2.0
2004	11.0	7.9	7.2	6.1	5.9	4.4	3.6	4.0	1.9
2005	11.0	8.4	7.3	6.6	6.1	4.2	3.5	4.0	1.9
2006	11.1	8.0	7.4	6.5	6.2	4.0	3.5	2.9	2.0
2007	11.5	7.9	7.7	6.7	6.1	3.7	3.5	2.7	2.0
2008	11.9	8.5	8.0	6.7	5.9	3.4	3.4	2.6	-

All rates per 100,000 population. A "-" indicates that the data are not available.

1. CLRD consists principally of bronchitis, emphysema, asthma, and chronic airways obstruction.
2. Beginning in 2006, the National Center for Health Statistics changed the ICD-10 codes to include only ICD-10 code I70.
3. Includes the alcohol-linked disorders represented by ICD-9 codes 291.0-291.9, 303, 305.0, 357.5, 425.5, 535.5 and 571.0-571.3, prior to 1999. For current ICD-10 components, see Table 6-6, footnotes 38-39.

NOTE: Beginning in 1999, causes of death were classified using the rubrics and methodology of the tenth revision of the International Classification of Disease (which supplanted the ninth revision). Final ICD-9/ICD-10 comparability ratios have been applied to all rates prior to 1999, except ALS and alcohol-induced deaths, where ratios were not calculated. See Appendix B.