

TABLE 6-53. Crude Death Rates¹ for Selected Leading Causes of Mortality, United States, 1995-2009²

Year	Total	Heart Disease	Cancer	CLRD	Cerebrovascular Disease	Unintentional Injuries	Alzheimer's Disease	Diabetes	Pneumonia & Influenza
1995	880.0	276.5	206.8	40.8	63.1	36.4	13.3	23.0	22.3
1996	872.5	272.4	205.3	41.6	63.3	36.7	13.4	23.7	22.3
1997	864.7	267.6	203.5	42.4	62.7	36.6	13.8	23.9	22.5
1998	864.2	263.7	202.1	43.1	58.9	37.1	13.8	24.4	23.7
1999	857.0	259.9	197.0	44.5	60.0	35.1	16.0	24.5	22.8
2000	854.0	252.6	196.5	43.4	59.6	34.8	17.6	24.6	23.2
2001	846.9	245.4	194.1	43.1	57.3	35.6	18.9	25.0	21.7
2002	847.3	241.7	193.2	43.3	56.4	37.0	20.4	25.4	22.8
2003	841.9	235.6	191.5	43.5	54.2	37.6	21.8	25.5	22.4
2004	816.5	222.2	188.6	41.5	51.1	38.1	22.5	24.9	20.3
2005	825.9	220.0	188.7	44.2	48.4	39.7	24.2	25.3	21.3
2006	810.4	211.0	187.0	41.6	45.8	40.6	24.2	24.2	18.8
2007	803.6	204.3	186.6	42.4	45.1	41.0	24.7	23.7	17.5
2008	813.0	202.9	186.0	46.4	44.1	40.1	27.1	23.2	18.5
2009	793.8	195.2	184.9	44.7	42.0	38.4	25.7	22.4	17.5

Year	Suicide	Hypertension	Alcohol ³	Parkinson's Disease	Homicide	HIV/AIDS	Congenital Anomalies	Arterio-sclerosis ⁴	ALS
1995	11.9	5.2	7.6	4.1	8.6	17.7	4.1	6.2	1.5
1996	11.6	5.5	7.3	4.5	7.8	12.7	4.0	6.1	1.6
1997	11.4	5.7	7.2	4.6	7.3	6.7	3.9	5.8	1.6
1998	11.3	5.9	7.1	4.9	6.6	5.4	3.9	5.5	1.6
1999	10.5	6.1	7.0	5.2	6.1	5.3	3.7	5.4	1.8
2000	10.4	6.4	7.0	5.6	6.0	5.1	3.8	5.1	1.9
2001	10.7	6.7	7.0	5.8	7.1	5.0	3.7	4.9	1.9
2002	11.0	7.0	7.0	5.9	6.1	4.9	3.7	4.8	2.0
2003	10.8	7.5	7.1	6.2	6.1	4.7	3.6	4.5	2.0
2004	11.0	7.9	7.2	6.1	5.9	4.4	3.6	4.0	1.9
2005	11.0	8.4	7.3	6.6	6.1	4.2	3.5	4.0	2.0
2006	11.1	8.0	7.4	6.5	6.2	4.0	3.5	2.9	2.0
2007	11.5	7.9	7.7	6.7	6.1	3.7	3.5	2.7	2.0
2008	11.9	8.5	8.0	6.7	5.9	3.4	3.4	2.6	2.0
2009	12.0	8.4	8.0	6.7	5.5	3.1	3.2	2.4	2.1

¹ All rates per 100,000 population.

² Most recent year for which final data are available.

³ See footnote for this cause in table 6-6. Prior to 1999, alcohol-induced deaths included ICD-9 codes 291, 303, 305.0, 357.5, 425.5, 535.5, and 571.0-571.3.

⁴ Beginning in 2006, the National Center for Health Statistics changed the ICD-10 codes for arteriosclerosis to include only ICD-10 code I70.

NOTE: Beginning in 1999, causes of death were classified using the rubrics and methodology of the tenth revision of the International Classification of Diseases (which supplanted the ninth revision). Final ICD-9/ICD-10 comparability ratios have been applied to all rates prior to 1999, except ALS and alcohol-induced deaths, where ratios were not calculated. See Appendix B.