

TABLE 6-57. Age-adjusted Death Rates for Selected Causes of Death, Oregon and United States Residents, 1996-2010¹

Year	Total			Cancer			Heart Disease		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1996	881.9	893.7	-1.3	208.8	208.6	0.1	230.6	281.4	-18.1
1997	864.0	877.5	-1.5	205.7	205.3	0.2	221.8	273.5	-18.9
1998	862.9	870.1	-0.8	207.9	202.5	2.7	210.7	267.2	-21.1
1999	845.3	875.6	-3.5	199.2	200.8	-0.8	208.0	266.5	-22.0
2000	826.9	869.0	-4.8	197.6	199.6	-1.0	197.5	257.6	-23.3
2001	835.9	851.6	-1.8	198.7	195.6	1.6	195.2	246.8	-20.9
2002	855.0	845.3	1.1	200.9	193.5	3.8	198.0	240.8	-17.8
2003	838.4	832.7	0.7	198.3	190.1	4.3	189.5	232.3	-18.4
2004	814.8	800.8	1.7	196.7	185.8	5.9	179.2	217.0	-17.4
2005	791.4	798.8	-0.9	189.4	183.8	3.0	169.5	211.1	-19.7
2006	784.5	776.5	1.0	185.7	180.7	2.8	162.6	200.2	-18.8
2007	771.6	760.2	1.5	184.7	178.4	3.5	159.7	190.9	-16.3
2008	772.8	758.3	1.9	182.8	175.3	4.3	154.5	186.5	-17.2
2009	739.7	741.1	-0.2	176.7	173.2	2.0	143.0	180.1	-20.6
2010	735.0	747.0	-1.6	177.9	172.8	2.9	139.7	179.1	-22.0

Year	Chronic Lower Resp. Disease			Cerebrovascular Disease			Unintentional Injuries		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1996	52.5	42.4	23.8	83.7	65.7	27.4	40.6	36.6	10.9
1997	50.8	42.8	18.7	80.8	64.3	25.7	39.9	36.4	9.6
1998	49.6	43.5	14.0	80.7	62.4	29.3	40.8	36.7	11.2
1999	50.4	45.4	11.0	80.3	61.6	30.4	33.9	35.3	-4.0
2000	47.8	44.2	8.1	70.8	60.9	16.3	34.6	34.9	-0.9
2001	48.7	43.6	11.7	71.4	57.7	23.7	35.4	35.5	-0.3
2002	50.9	43.5	17.0	71.7	56.2	27.6	38.4	36.9	4.1
2003	49.8	43.3	15.0	68.5	53.5	28.0	38.3	37.3	2.7
2004	48.1	41.1	17.0	61.9	50.0	23.8	38.8	37.7	2.9
2005	47.8	43.2	10.6	57.3	46.6	23.0	37.6	39.1	-3.8
2006	46.8	40.5	15.6	48.8	43.6	11.9	40.7	39.8	2.3
2007	47.5	40.8	16.4	44.5	42.2	5.5	41.7	40.0	4.3
2008	48.2	44.0	9.5	45.6	40.7	12.0	42.4	38.8	9.3
2009	46.4	42.3	9.6	44.0	38.9	13.2	38.8	37.3	3.9
2010	46.5	42.2	10.2	40.5	39.1	3.6	37.8	38.0	-0.6

¹ Most recent year for which final US data are available.

NOTE: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in Tables 6-54 and 6-55, all Oregon data are from state mortality files. Consequently, the rates and percentage differences shown here will vary from those in Tables 6-54 and 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. National comparability ratios have been applied to rates prior to 1999 to control for coding changes resulting from the implementation of ICD-10 when it supplanted ICD-9 in 1999. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease vs. Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the State's query program: when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information.

TABLE 6-57. Age-adjusted Death Rates for Selected Causes of Death, Oregon and United States Residents, 1996-2010¹ — Continued

Year	Alzheimer's Disease			Diabetes Mellitus			Suicide		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1996	20.6	13.4	53.7	23.0	24.3	-5.3	16.7	11.5	45.2
1997	19.8	13.8	43.5	24.9	24.2	2.9	16.7	11.2	49.1
1998	19.0	13.6	39.7	26.0	24.6	5.7	17.2	11.1	55.0
1999	24.7	16.5	49.7	24.7	25.0	-1.2	14.9	10.5	41.9
2000	24.8	18.1	37.0	23.8	25.0	-4.8	14.3	10.4	37.5
2001	28.1	19.0	47.9	28.8	25.2	14.3	14.9	10.7	39.3
2002	30.3	20.2	50.0	28.6	25.4	12.6	14.5	10.9	33.0
2003	30.6	21.4	43.0	28.1	25.3	11.1	16.3	10.8	50.9
2004	33.4	21.8	53.2	29.0	24.5	18.4	15.2	10.9	39.4
2005	30.4	22.9	32.8	29.3	24.6	19.1	14.9	10.9	36.7
2006	29.5	22.6	30.5	28.9	23.3	24.0	15.1	10.9	38.5
2007	28.0	22.7	23.3	27.9	22.5	24.0	15.6	11.3	38.1
2008	30.5	24.4	25.0	24.8	21.8	13.8	14.7	11.6	26.7
2009	27.7	23.5	17.8	25.3	20.9	20.9	16.1	11.8	36.2
2010	28.7	25.1	14.3	24.2	20.8	16.3	17.1	12.1	41.4

Year	Alcohol-Induced			Hypertension			Flu & Pneumonia		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1996	12.8	8.2	56.1	6.5	5.7	14.0	20.1	22.9	-12.2
1997	11.5	7.9	45.6	7.7	5.8	32.8	19.0	23.2	-18.1
1998	11.0	7.8	41.0	6.6	6.0	10.0	20.7	24.1	-14.1
1999	8.9	7.1	25.4	7.0	6.2	12.9	19.5	23.5	-17.0
2000	10.8	7.0	54.3	6.2	6.5	-4.6	17.5	23.7	-26.2
2001	12.2	7.0	74.3	8.6	6.8	26.5	15.7	21.9	-28.3
2002	12.3	6.9	78.3	9.6	7.0	37.1	17.9	22.6	-20.8
2003	14.2	7.0	102.9	9.3	7.4	25.7	17.0	22.0	-22.7
2004	13.8	7.0	97.1	9.5	7.7	23.4	14.7	19.8	-25.8
2005	13.7	7.0	95.7	10.6	8.0	32.5	15.1	20.3	-25.6
2006	11.7	7.0	67.1	8.9	7.5	18.7	12.8	17.8	-28.1
2007	13.1	7.3	79.5	8.6	7.4	16.2	11.4	16.2	-29.6
2008	12.9	7.4	74.3	9.5	7.7	23.4	12.3	16.9	-27.2
2009	13.4	7.4	81.4	9.5	7.7	23.1	12.0	16.2	-26.2
2010	13.0	7.6	71.2	9.8	8.0	23.1	9.3	15.1	-38.4

¹ Most recent year for which final US data are available.

NOTE: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in Tables 6-54 and 6-55, all Oregon data are from state mortality files. Consequently, the rates and percentage differences shown here will vary from those in Tables 6-54 and 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. National comparability ratios have been applied to rates prior to 1999 to control for coding changes resulting from the implementation of ICD-10 when it supplanted ICD-9 in 1999. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease vs. Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the State's query program: when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information.

TABLE 6-57. Age-adjusted Death Rates for Selected Causes of Death, Oregon and United States Residents, 1996-2010¹ — Continued

Year	Parkinson's Disease			Viral Hepatitis			Homicide		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1996	7.2	4.6	56.5	1.1	1.0	10.0	4.5	7.5	-40.0
1997	6.4	4.7	36.2	1.4	1.1	27.3	3.9	7.0	-44.3
1998	8.0	4.9	63.3	1.6	1.3	23.1	4.1	6.4	-35.9
1999	7.3	5.4	35.2	1.3	1.8	-27.8	3.3	6.0	-45.0
2000	7.7	5.7	35.1	2.2	1.9	15.8	2.7	5.9	-54.2
2001	8.0	5.8	37.9	2.5	2.0	25.0	3.1	7.1	-56.3
2002	8.3	5.9	40.7	3.5	2.0	75.0	3.1	6.1	-49.2
2003	8.4	6.2	35.5	2.6	1.8	44.4	2.5	6.0	-58.3
2004	8.6	6.1	41.0	2.9	1.8	61.1	3.1	5.9	-47.5
2005	7.7	6.4	20.3	2.3	1.8	27.8	2.9	6.1	-52.5
2006	8.7	6.3	38.1	2.2	2.3	-4.3	3.0	6.2	-51.6
2007	8.2	6.4	28.1	4.2	2.3	82.6	2.1	6.1	-65.6
2008	8.7	6.4	35.9	3.8	2.3	65.2	2.6	5.9	-55.9
2009	8.3	6.4	29.7	3.9	2.2	77.1	2.6	5.5	-53.3
2010	8.5	6.8	25.6	3.8	2.1	79.8	2.9	5.3	-45.5

Year	Amyotrophic Lateral Sclerosis			Arteriosclerosis			HIV/AIDS		
	Oregon	US	% Diff	Oregon	US	% Diff	Oregon	US	% Diff
1996	2.0	1.6	25.0	7.5	6.4	17.2	7.6	11.9	-36.1
1997	2.3	1.6	43.8	6.9	6.0	15.0	3.2	6.2	-48.4
1998	2.2	1.6	37.5	6.5	5.6	16.1	2.3	4.9	-53.1
1999	2.2	1.9	15.8	5.6	5.5	1.8	2.2	5.3	-58.5
2000	2.7	2.0	35.0	6.4	5.2	23.1	1.8	5.2	-65.4
2001	2.6	1.9	36.8	5.3	5.0	6.0	1.9	5.0	-62.0
2002	3.0	2.0	50.0	5.7	4.7	21.3	2.5	4.9	-49.0
2003	3.1	2.0	55.0	5.5	4.4	25.0	2.5	4.7	-46.8
2004	2.9	1.9	52.6	4.6	3.9	17.9	1.8	4.5	-60.0
2005	2.8	1.9	47.4	4.8	3.8	26.3	1.5	4.2	-64.3
2006	2.9	1.9	52.6	2.8	2.7	3.7	1.4	4.0	-65.0
2007	2.3	1.9	21.1	3.0	2.5	20.0	1.5	3.7	-59.5
2008	3.0	1.9	57.9	2.2	2.3	-4.3	1.0	3.3	-69.7
2009	2.7	1.9	39.8	1.8	2.2	-19.0	1.1	3.0	-62.7
2010	2.8	2.0	38.9	1.6	2.2	-29.1	1.2	2.6	-55.1

¹ Most recent year for which final US data are available.

NOTE: US age-adjusted death rates are from compressed mortality files available at the federal Centers for Disease Control and Prevention's WONDER online database. Unlike the data shown in Tables 6-54 and 6-55, all Oregon data are from state mortality files. Consequently, the rates and percentage differences shown here will vary from those in Tables 6-54 and 6-55 due to different file closure dates, different population estimate methodologies, and incorporation of physician query results. National comparability ratios have been applied to rates prior to 1999 to control for coding changes resulting from the implementation of ICD-10 when it supplanted ICD-9 in 1999. Because the ratios are based on national data, discontinuities may occur when Oregon physicians reported causes of death differently than their national counterparts (e.g., Alzheimer's disease vs. Alzheimer's dementia). Some differences between Oregon and US rates (e.g., alcohol-induced deaths) result, at least in part, from the State's query program: when death certificates are incomplete, letters are sent to physicians/certifiers requesting additional information.