

TABLE 6-13. Years of potential life lost before age 75 from the leading causes of death, by year, Oregon residents, 1999-2013

Year	Total	Cancer	Unintended injury	Heart disease	Suicide	Alcohol-induced ¹	Perinatal conditions	Diabetes	CLRD ²
1999	207,575	48,693	29,370	30,546	13,915	5,645	8,396	5,546	5,878
2000	206,973	49,688	31,398	27,487	14,317	6,778	7,856	4,839	5,487
2001	211,233	51,244	30,249	27,225	15,023	7,821	8,396	5,852	5,567
2002	222,274	52,637	31,185	28,489	14,455	8,125	8,966	5,929	5,802
2003	225,545	50,810	34,383	28,869	15,585	10,033	8,591	7,237	6,493
2004	221,453	50,892	34,830	26,449	15,294	9,877	8,396	7,497	5,848
2005	224,868	53,166	31,845	26,721	14,874	9,553	10,131	7,585	6,543
2006	231,592	52,025	36,529	26,871	16,158	9,082	9,067	7,590	6,807
2007	234,443	51,747	36,820	27,845	16,266	10,168	10,311	7,551	7,307
2008	231,750	51,479	38,621	27,793	16,342	10,362	8,994	6,621	7,598
2009	230,153	53,568	34,029	25,605	17,158	10,686	8,323	7,530	7,341
2010	224,366	54,941	30,199	23,929	17,963	10,666	7,891	7,292	7,799
2011	230,525	55,353	33,117	24,368	18,023	11,984	8,201	7,831	7,604
2012	228,909	54,352	31,236	24,889	19,481	11,856	8,473	7,273	7,141
2013	233,367	53,926	30,610	24,786	19,119	12,867	9,188	7,665	8,121

Year	Congenital anomalies	Cerebro-vascular disease	Homicide ³	Viral hepatitis	Undetermined external causes	Sudden infant death syndrome	Pneumonia & influenza	HIV disease	Septicemia
1999	7,846	5,629	4,804	989	2,146	1,939	1,519	2,420	1,656
2000	6,556	5,276	3,798	1,713	2,040	3,802	1,301	2,040	1,446
2001	6,844	6,011	3,887	1,681	2,663	2,162	1,873	2,050	1,240
2002	7,439	6,012	4,728	2,560	3,592	2,310	2,344	2,691	1,423
2003	6,313	6,108	3,522	2,050	3,575	1,714	1,985	2,675	1,309
2004	6,720	6,221	4,502	2,105	3,284	1,416	1,671	1,902	1,481
2005	5,695	6,274	4,078	1,717	3,370	1,491	2,421	1,729	1,658
2006	6,918	5,737	4,429	1,817	3,390	2,236	1,578	1,478	1,429
2007	6,293	6,339	3,147	3,536	3,691	2,833	1,684	1,518	1,709
2008	6,271	5,135	3,949	2,860	2,693	1,492	2,236	1,045	1,839
2009	4,264	5,714	3,684	3,276	3,004	2,163	3,822	1,076	2,096
2010	5,688	5,206	4,080	3,197	3,432	2,385	1,760	1,130	1,660
2011	5,831	5,709	4,235	3,177	2,437	2,087	1,786	859	1,581
2012	5,405	5,171	4,159	2,597	2,379	1,865	1,482	1,359	1,253
2013	5,607	5,302	3,211	3,858	2,316	1,715	1,915	1,234	1,403

¹ See Table 6-6, footnotes 38-39, for a list of included conditions and their ICD codes. Prior to 1999, figures do not include deaths due to alcohol poisoning.

² Chronic lower respiratory disease.

³ Excludes legal intervention.