

TABLE 6-54. Crude death rates¹ for selected leading causes of mortality, United States, 1999-2013²

Year	Total	Heart disease	Cancer	CLRD	Unintentional injuries	Cerebro-vascular disease	Alzheimer's disease	Diabetes	Pneumonia & influenza
1999	857.0	259.9	197.0	44.5	35.1	60.0	16.0	24.5	22.8
2000	854.0	252.6	196.5	43.4	34.8	59.6	17.6	24.6	23.2
2001	846.9	245.4	194.1	43.1	35.6	57.3	18.9	25.0	21.7
2002	847.3	241.7	193.2	43.3	37.0	56.4	20.4	25.4	22.8
2003	841.9	235.6	191.5	43.5	37.6	54.2	21.8	25.5	22.4
2004	816.5	222.2	188.6	41.5	38.1	51.1	22.5	24.9	20.3
2005	825.9	220.0	188.7	44.2	39.7	48.4	24.2	25.3	21.3
2006	810.4	211.0	187.0	41.6	40.6	45.8	24.2	24.2	18.8
2007	803.6	204.3	186.6	42.4	41.0	45.1	24.7	23.7	17.5
2008	813.0	202.9	186.0	46.4	40.1	44.1	27.1	23.2	18.5
2009	793.8	195.2	184.9	44.7	38.4	42.0	25.7	22.4	17.5
2010	799.5	193.6	186.2	44.7	39.1	41.9	27.0	22.4	16.2
2011	807.3	191.5	185.1	45.9	40.6	41.4	27.3	23.7	17.3
2012	810.2	191.0	185.6	45.7	40.7	40.9	26.6	23.6	16.1
2013	821.5	193.3	185.0	47.2	41.3	40.8	26.8	23.9	18.0

Year	Suicide	Hypertension	Alcohol ³	Parkinson's disease	Homicide	Congenital anomalies	HIV/AIDS	ALS	Arterio-sclerosis ⁴
1999	10.5	6.1	7.0	5.2	6.1	3.7	5.3	1.8	5.4
2000	10.4	6.4	7.0	5.6	6.0	3.8	5.1	1.9	5.1
2001	10.7	6.7	7.0	5.8	7.1	3.7	5.0	1.9	4.9
2002	11.0	7.0	7.0	5.9	6.1	3.7	4.9	2.0	4.8
2003	10.8	7.5	7.1	6.2	6.1	3.6	4.7	2.0	4.5
2004	11.0	7.9	7.2	6.1	5.9	3.6	4.4	1.9	4.0
2005	11.0	8.4	7.3	6.6	6.1	3.5	4.2	2.0	4.0
2006	11.1	8.0	7.4	6.5	6.2	3.5	4.0	2.0	2.9
2007	11.5	7.9	7.7	6.7	6.1	3.5	3.7	2.0	2.7
2008	11.9	8.5	8.0	6.7	5.9	3.4	3.4	2.0	2.6
2009	12.0	8.4	8.0	6.7	5.5	3.2	3.1	2.1	2.4
2010	12.4	8.6	8.3	7.1	5.3	3.1	2.7	2.2	2.3
2011	12.7	8.9	8.6	7.4	5.2	3.1	2.5	2.2	2.2
2012	12.9	9.3	8.8	7.6	5.3	3.1	2.3	2.3	2.2
2013	13.0	9.7	9.2	8.0	5.1	3.0	2.2	2.2	2.1

¹ All rates per 100,000 population.

² Most recent year for which final data are available.

³ See footnote for this cause in table 6-6. Prior to 1999, alcohol-induced deaths included ICD-9 codes 291, 303, 305.0, 357.5, 425.5, 535.5, and 571.0-571.3.

⁴ Beginning in 2006, the National Center for Health Statistics changed the ICD-10 codes for arteriosclerosis to include only ICD-10 code I70.

NOTE: Beginning in 1999, causes of death were classified using the rubrics and methodology of the tenth revision of the International Classification of Diseases (which supplanted the ninth revision). Final ICD-9/ICD-10 comparability ratios have been applied to all rates prior to 1999, except ALS and alcohol-induced deaths, where ratios were not calculated. See Appendix B.