

TABLE 6-20. Tobacco-linked deaths by cause of death, Oregon residents, 2015

Selected causes of death (and their ICD-10 codes)	Total	Linked ¹		Not linked		Unknown	
		Number	Percent	Number	Percent	Number	Percent
Total ²	35,709	7,670	21.5	19,517	54.7	8,522	23.9
Malignant neoplasms	3,752	1,895	50.5	1,142	30.4	715	19.1
Oral cavity, lip, pharynx (C00.0-C14.8)	166	83	50.0	44	26.5	39	23.5
Esophagus (C15)	222	81	36.5	69	31.1	72	32.4
Stomach (C16)	106	19	17.9	57	53.8	30	28.3
Pancreas (C25)	638	68	10.7	415	65.0	155	24.3
Larynx (C32)	24	17	70.8	4	16.7	3	12.5
Lung, bronchi, and trachea (C33-C34)	1,956	1,479	75.6	202	10.3	275	14.1
Cervix uteri (C53)	53	8	15.1	38	71.7	7	13.2
Kidney, other urinary tract (C64-C65)	200	33	16.5	117	58.5	50	25.0
Urinary bladder (C67)	260	98	37.7	99	38.1	63	24.2
Acute myeloid leukemia (C92.0)	127	9	7.1	97	76.4	21	16.5
Cardiovascular disease	8,774	1,933	22.0	4,276	48.7	2,565	29.2
Ischemic heart disease (I20-I25)	3,439	1,081	31.4	1,400	40.7	958	27.9
Other heart disease (I00-I09, I26-I51)	3,133	462	14.7	1,753	56.0	918	29.3
Cerebrovascular disease (I60-I69)	1,869	262	14.0	1,005	53.8	602	32.2
Atherosclerosis (I70)	47	14	29.8	25	53.2	8	17.0
Aortic aneurysm (I71)	152	59	38.8	44	28.9	49	32.2
Other arterial disease (I72-I78)	134	55	41.0	49	36.6	30	22.4
Respiratory diseases	2,489	1,674	67.3	421	16.9	394	15.8
Pneumonia and influenza (J09-J18)	453	56	12.4	265	58.5	132	29.1
Bronchitis and emphysema (J40-J43)	148	135	91.2	7	4.7	6	4.1
Other chronic airways obstruction (J44)	1,888	1,483	78.5	149	7.9	256	13.6
Perinatal conditions ³	71	—	—	66	93.0	5	7.0
Selected perinatal conditions ⁴	48	—	—	44	91.7	4	8.3
Sudden infant death syndrome (R95)	23	—	—	22	95.7	1	4.3
Other causes	20,623	2,168	10.5	13,612	66.0	4,843	23.5

[—] Quantity is zero.

¹ The Oregon death certificate provides four possible answers to the question, "Did tobacco use contribute to death?": yes, probably, no and unknown. The linked category includes deaths listed as yes or probably, or if a contributing cause is F17.

² The causes of death shown in this table are those linked to tobacco use by the federal Centers for Disease Control and Prevention (CDC. Annual Smoking-Attributable Mortality, Years of Potential Life Lost, and Productivity Losses -- United States, 1997-2001. MMWR 2005; 54:625-628.).

³ The number of infant deaths resulting from exposure to tobacco combustion products is underreported.

⁴ The category includes the following conditions: other disorders related to short gestation and low birthweight (P07), respiratory distress of newborn (P22), congenital pneumonia (P23), neonatal aspiration syndromes (P24), and other respiratory conditions originating in the perinatal period (P25-P28).