

**TABLE 7-8. Early neonatal deaths<sup>1</sup> by weeks of gestation and weight, Oregon residents, birth cohort 2016**

Birthweight (in grams)	Total	Weeks of gestation <sup>2</sup>									
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+	N.S.
Total <sup>3</sup> .....	119	11	47	20	11	9	1	16	3	–	1
001-349 .....	21	11	10	–	–	–	–	–	–	–	–
350-499 .....	22	–	19	3	–	–	–	–	–	–	–
<500 .....	43	11	29	3	–	–	–	–	–	–	–
500-749 .....	28	–	18	8	1	1	–	–	–	–	–
750-999 .....	7	–	–	6	–	–	–	–	–	–	1
1000-1249 .....	2	–	–	–	2	–	–	–	–	–	–
1250-1499 .....	4	–	–	1	1	2	–	–	–	–	–
1500-1999 .....	13	–	–	1	5	3	–	4	–	–	–
2000-2499 .....	3	–	–	–	2	–	1	–	–	–	–
<2500 .....	100	11	47	19	11	6	1	4	–	–	1
2500+ .....	18	–	–	–	–	3	–	12	3	–	–
2500-2999 .....	5	–	–	–	–	1	–	4	–	–	–
3000-3499 .....	8	–	–	–	–	2	–	5	1	–	–
3500-3999 .....	4	–	–	–	–	–	–	3	1	–	–
4000-4499 .....	–	–	–	–	–	–	–	–	–	–	–
4500+ .....	1	–	–	–	–	–	–	–	1	–	–

– Quantity is zero.

<sup>1</sup> Early neonatal deaths occur through day six after birth.

<sup>2</sup> Clinical estimate of gestation. If clinical estimate is unknown, gestation is calculated from reported date of last menses.

<sup>3</sup> Total includes reports with unknown weight.