

TABLE 7-9. Late neonatal deaths¹ by weeks of gestation and weight, Oregon residents, birth cohort 2016

Birthweight (in grams)	Total	Weeks of gestation ²								
		<20	20-23	24-27	28-31	32-35	36	37-39	40	41+
Total	32	–	1	13	1	3	1	8	4	1
001-349	–	–	–	–	–	–	–	–	–	–
350-499	–	–	–	–	–	–	–	–	–	–
<500	–	–	–	–	–	–	–	–	–	–
500-749	8	–	1	7	–	–	–	–	–	–
750-999	5	–	–	4	1	–	–	–	–	–
1000-1249	1	–	–	1	–	–	–	–	–	–
1250-1499	–	–	–	–	–	–	–	–	–	–
1500-1999	3	–	–	–	–	1	1	–	1	–
2000-2499	5	–	–	1	–	2	–	2	–	–
<2500	22	–	1	13	1	3	1	2	1	–
2500+	10	–	–	–	–	–	–	6	3	1
2500-2999	3	–	–	–	–	–	–	2	1	–
3000-3499	5	–	–	–	–	–	–	4	1	–
3500-3999	2	–	–	–	–	–	–	–	1	1
4000-4499	–	–	–	–	–	–	–	–	–	–
4500+	–	–	–	–	–	–	–	–	–	–

– Quantity is zero.

¹ Late neonatal deaths occur from day seven through 27 after birth.

² Clinical estimate of gestation. If clinical estimate is unknown, gestation is calculated from reported date of last menses.